IN THE CLAIMS:

Please rewrite claim 1 and cancel claims 10 to 12 and 27 to 34 without prejudice or disclaimer as follows:

1. (currently amended) A method to conduct an experiment, comprising steps of:

selecting factors for the experiment;

estimating interactions among levels of the factors

assigning a probability value of positive interactions for each of the estimated interactions;

effecting a combinatorial high throughput screening (CHTS) method on an experimental space representing the levels, wherein the CHTS method comprises defining a first experimental space by structuring the levels according to a Latin Square strategy; and

adjusting the probabilities for each interaction according to results of the CHTS method.

- 2. (original) The method of claim 1, comprising assigning a high probability value, medium probability value or low probability value of each positive interaction for each of the estimated interactions is assigned by a client or investigator.
- 3. (original) The method of claim 1, wherein a high probability value, medium probability value or low probability value of each positive interaction for each of the estimated interactions.
- 4. (original) The method of claim 1, wherein an investigator and a client who benefits from results from the CHTS experiment in concert determine a probability value to be assigned.

- 5. (original) The method of claim 1, comprising assigning values to represent a high probability value, medium probability value and low probability value of each positive interaction for each of the estimated interactions.
- 6. (original) The method of claim 1, comprising assigning 0.6 to about 0.99 value as a high probability value, about 0.2 to about 0.59 value as a medium probability value and about 0.01 to about 0.19 value as a low probability value.
- 7. (original) The method of claim 1, comprising assigning 0.7 to about 0.9 value as a high probability value, about 0.2 to about 0.5 value as a medium probability value and about .05 to about 0.15 value as a low probability value.
- 8. (original) The method of claim 1, further comprising repeating a CHTS method step and an adjusting probabilities step until a best set of levels is selected.
- 9. (original) The method of claim 1, comprising constructing an adjustable definitional model to represent the estimated interactions and assigned probabilities.
 - 10. (canceled)
 - 11. (canceled)
 - 12. (canceled)
- 13. (original) The method of claim 1, wherein the CHTS experiment comprises steps of;

preparing a plurality of reagent compositions;

formulating a combinatorial library of reactants from said plurality of reagent compositions;

effecting parallel reaction of the library to produce products; and evaluating the products to select a lead from the library of reactants.

- 14. (original) The method of claim 1, wherein conducting the CHTS experiment comprises providing a reactor plate comprising a substrate with an array of reaction cells containing at least one reactant according to an input factor level and reacting the reactant in parallel with other reactants
- 15. (original) The method of claim 1, wherein the CHTS comprises effecting parallel chemical reactions of an array of reactants defined as the experimental space.
- 16. (original) The method of claim 1, wherein the CHTS comprises effecting parallel chemical reactions on a micro scale on reactants defined as the experimental space.
- 17. (original) The method of claim 1, wherein the CHTS comprises an iteration of steps of simultaneously reacting a multiplicity of tagged reactants and identifying a multiplicity of tagged products of the reaction and evaluating the identified products after completion of a single or repeated iteration.
- 18. (original) The method of claim 1, wherein the experimental space factors comprise reactants, catalysts and conditions and the CHTS comprises
- (A) (a) reacting a reactant selected from the experimental space under a selected set of catalysts or reaction conditions; and (b) evaluating a set of results of the reacting step; and
- (B) reiterating step (A) wherein a selected experimental space selected for a step (a) is chosen as a result of an evaluating step (b) of a preceding iteration of step (A).
- 19. (original) The method of claim 16, wherein the evaluating step (b) comprises identifying relationships between factor levels of the candidate chemical reaction space; and determining the chemical experimental space according to a full factorial design for the next iteration.
- 20. (original) The method of claim 16, comprising reiterating (A) until a best set of factor levels of the chemical experimental space is selected.

- 21. (original) The method of claim 1, wherein the factors include a catalyst system comprising a Group VIII B metal.
- 22. (original) The method of claim 1, wherein the factors include a catalyst system comprising palladium.
- 23. (original) The method of claim 1, wherein the factors include a catalyst system comprising a halide composition.
- 24. (original) The method of claim 1, wherein the factors include an inorganic co-catalyst.
- 25. (original) The method of claim 1, wherein the factors include a catalyst system includes a combination of inorganic co-catalysts.
- 26. (original) The method of claim 1, wherein the factors comprise a reactant or catalyst at least partially embodied in a liquid and effecting the CHTS method comprises contacting the reactant or catalyst with an additional reactant at least partially embodied in a gas, wherein the liquid forms a film having a thickness sufficient to allow a reaction rate that is essentially independent of a mass transfer rate of additional reactant into the liquid to synthesize products that comprise the results.
 - 27. (canceled)
 - 28. (canceled)
 - 29. (canceled)
 - 30. (canceled)
 - 31. (canceled)
 - 32. (canceled)
 - 33. (canceled)
 - 34. (canceled)